
GRN008978 Greenacle Poultry Dosage and Expectations for Laying Hens

Please see the treatment for Laying Hens and expected results:

DOSAGE:

You can use Greenacle Poultry liquid or Greenacle Poultry granulate.

Greenacle Poultry liquid: 1 liter of additive to 2,500 liters of water. Please always add additive to water and not vice versa. The prepared solution must be used within 72 hours. After that the effects will be reduced or completely lost. If the solution is older than 72 hours, please prepare fresh solution for the treatment.

Greenacle Poultry granulate: 600gr of granulate in 1,000 kg of food. Please make sure the granulate is well mixed into the food. The solution can be used for 2-3 months.

IMPORTANT: It is necessary to immediately reduce food by 10% so that the chickens do not grow. If you observe that chickens are growing (getting fat) reduce the food even more.

EXPECTED RESULTS:

- Egg production is increased;
- The hens start laying eggs earlier and will lay eggs for longer time;
- Broken eggs are reduced as the shell is stronger;
- Eggs are of the better quality, more beta-carotene, less cholesterol, bigger yolk, a better ratio between omega 3 and 6;
- The mortality of hens is reduced, especially when in the 4th quarter of laying;
- Ammonium in the farm decrease for about 50%;
- Improved food conversion in the birds digestive system;
- Food savings from the beginning of the treatment.